

A Property of Kähler-Ricci Solitons on Complete Complex Surfaces

Bing-Long Chen and Xi-Ping Zhu

Department of Mathematics,
Zhongshan University,
Guangzhou 510275, P. R. China

1. Introduction

This paper is concerned with Ricci flow on complete Kähler manifolds. Let M be a complete complex manifolds with Kähler metric $g_{\alpha\bar{\beta}}$. The Ricci flow is the following evolution equation on the metric

$$\begin{cases} \frac{\partial g_{\alpha\bar{\beta}}(x, t)}{\partial t} = -R_{\alpha\bar{\beta}}(x, t) & x \in M \quad t > 0, \\ g_{\alpha\bar{\beta}}(x, 0) = g_{\alpha\bar{\beta}}(x) & x \in M, \end{cases} \quad (1.1)$$

where $R_{\alpha\bar{\beta}}(x, t)$ denotes the Ricci curvature tensor of the metric $g_{\alpha\bar{\beta}}(x, t)$.

One of the main problems in differential geometry is to find canonical structure on manifolds. The Ricci flow introduced by Hamilton [8] is an useful tool to approach such problems. For examples, Hamilton [10] and Chow [7] used the convergence of the Ricci flow to characterize the complex structures on compact Riemann surfaces, Hamilton [8] used the Ricci flow to classify compact three-manifolds with positive Ricci curvature, and the authors [4] recently used the Ricci flow to get steinness for a class of complete noncompact Kähler manifolds. By a direct computation one can see that the scalar curvature $R(x, t)$ of $g_{\alpha\bar{\beta}}(x, t)$ satisfies the equation

$$\frac{\partial}{\partial t} R = \Delta R + 2 |R_{\alpha\bar{\beta}}|^2.$$

This is a nonlinear heat equation with superlinear growth. It is clear that the scalar curvature must generally blow up in finite time. In the other words

the Ricci flow will develop singularities in finite time. Thus it is important to consider what kind of singularities might form.

In [12], Hamilton divided the solutions of the Ricci flow into three types and showed that one can dilate the solutions around the singularities to get in limit the models of the corresponding type. Then Cao [2] showed that if in addition the limits of Type II or Type III have nonnegative holomorphic bisectional curvature, the singularity models are the solutions to (1.1) which move by one-parameter group of biholomorphisms or also expand by a factor at the same time.

A solution to the Ricci flow (1.1) which moves by a one-parameter group of biholomorphisms is called a Kähler-Ricci soliton. The equation for a Kähler metric to move by a biholomorphism in the direction of a holomorphic vector field V is that the Ricci term $R_{\alpha\bar{\beta}}$ is the Lie derivative of the metric $g_{\alpha\bar{\beta}}$ in the direction of the vector field V ; thus

$$\begin{cases} R_{\alpha\bar{\beta}} = g_{\alpha\bar{\gamma}} \nabla_{\bar{\beta}} V^{\bar{\gamma}} + g_{\gamma\bar{\beta}} \nabla_{\alpha} V^{\gamma}, \\ \nabla_{\alpha} V^{\bar{\beta}} = \nabla_{\bar{\beta}} V^{\alpha} = 0. \end{cases} \quad (1.2)$$

From [12], any Kähler-Ricci soliton must be noncompact. Recall that a complete noncompact complex n -dimensional Kähler manifold $(M, g_{\alpha\bar{\beta}})$ is of maximal volume growth if there exist some fixed point $x_0 \in M$ and some positive constant C_0 such that

$$\text{Vol}(B(x_0, r)) \geq C_0 r^{2n} \quad \text{for all} \quad 0 \leq r < +\infty.$$

where $\text{Vol}(B(x_0, r))$ denotes the volume of the geodesic ball $B(x_0, r)$ centered at x_0 with radius r . The main result of this paper is the following property for Kähler-Ricci solitons.

Theorem Let $(M, g_{\alpha\bar{\beta}})$ be a nonflat Kähler-Ricci soliton on a complex two-dimensional Kähler manifold with bounded and nonnegative holomorphic bisectional curvature. Then $(M, g_{\alpha\bar{\beta}})$ can not be of maximal volume growth.

We conjecture this theorem holds for all dimensions. In the forthcoming paper [5], we will use this result to obtain a uniformization theorem for complete noncompact Kähler surfaces with positive holomorphic bisectional curvature.

This paper contains three sections. In the next section we will show that the curvature operator of the Kähler-Ricci soliton is nonnegative definite everywhere. In the last section we will use a blow down argument to prove the theorem.

We are grateful to H.D. Cao for many helpful discussions. This work was partially supported by the Foundation for Outstanding Young Scholars of China.

2. Nonnegativity of Curvature Operator

The solution to the Ricci flow (1.1) corresponding to the Kähler-Ricci soliton exists for $-\infty < t < +\infty$ and is obtained by flowing along the holomorphic vector field V in (1.2). The following lemma shows that the nonnegativity of holomorphic bisectional curvature of the Kähler-Ricci soliton on a complex surface implies the nonnegativity of its curvature operator.

Lemma Let $g_{\alpha\bar{\beta}}(x, t)$ be a complete solution to the Ricci flow (1.1) on a complex two-dimensional Kähler manifold M for all $t \in (-\infty, +\infty)$. Suppose its holomorphic bisectional curvature is bounded and nonnegative everywhere. Then the curvature operator of the metric $g_{\alpha\bar{\beta}}(x, t)$ is nonnegative definite everywhere on $M \times (-\infty, +\infty)$.

Proof. Choose a local orthonormal coframe $\{\omega_1, \omega_2, \omega_3, \omega_4\}$ on an open set $U \subset M$ so that $\omega_1 + \sqrt{-1}\omega_2$ and $\omega_3 + \sqrt{-1}\omega_4$ are $(1, 0)$ -forms over U . Then the self-dual forms

$$\varphi_1 = \omega_1 \wedge \omega_2 + \omega_3 \wedge \omega_4, \quad \varphi_2 = \omega_2 \wedge \omega_3 + \omega_1 \wedge \omega_4, \quad \varphi_3 = \omega_3 \wedge \omega_1 + \omega_2 \wedge \omega_4$$

and the anti-self-dual forms

$$\psi_1 = \omega_1 \wedge \omega_2 - \omega_3 \wedge \omega_4, \quad \psi_2 = \omega_2 \wedge \omega_3 - \omega_1 \wedge \omega_4, \quad \psi_3 = \omega_3 \wedge \omega_1 - \omega_2 \wedge \omega_4$$

form a basis of the space of 2-forms over U . In particular, $\varphi_1, \psi_1, \psi_2$ and ψ_3 give a basis for the space of $(1, 1)$ -forms over U .

On a complex two-dimensional Kähler manifold, it is well known that its curvature operator has image in the holonomy algebra $u(2) (\subset so(4))$ spanned by $(1, 1)$ -forms. Thus, the curvature operator R_m in the basis $\{\varphi_1, \varphi_2, \varphi_3, \psi_1, \psi_2, \psi_3\}$ has the following form,

$$R_m = \begin{pmatrix} a & 0 & 0 & b_1 & b_2 & b_3 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ b_1 & 0 & 0 & & & \\ b_2 & 0 & 0 & & A & \\ b_3 & 0 & 0 & & & \end{pmatrix}$$

where A is a 3×3 symmetric matrix.

Let V be a real tangent vector of the complex surface M . Denote by J the complex structure. It is clear that the complex 2-plane $V \wedge JV$ is dual to $(1, 1)$ -form $u\varphi_1 + v_1\psi_1 + v_2\psi_2 + v_3\psi_3$ satisfying the decomposability condition $u^2 = v_1^2 + v_2^2 + v_3^2$. Then after normalizing u to 1 by scaling, we see that the holomorphic bisectional curvature is nonnegative if and only if

$$a + b \cdot v + b \cdot w + {}^t v A w \geq 0, \quad (2.1)$$

for any unit vectors $v = (v_1, v_2, v_3)$ and $w = (w_1, w_2, w_3)$ in $\mathbf{R}^3$, where b is the vector (b_1, b_2, b_3) in R_m .

Denote by $a_1 \leq a_2 \leq a_3$ the eigenvalues of A . Recall that $\text{tr } A = a$ by the Bianchi identity, so if we choose v to be the eigenvector of A with eigenvalue a_3 and choose $w = -v$, (2.1) gives

$$a_1 + a_2 \geq 0. \quad (2.2)$$

In particular, we have $a_2 \geq 0$.

To proceed further, we need to adapt the Hamilton's maximum principle for tensors. Let

$$(a_i)_{\min}(t) = \inf_{x \in M} a_i(x, t), \quad i = 1, 2, 3$$

and

$$K = \sup_{(x,t) \in M \times (-\infty, +\infty)} |Rm(x, t)|.$$

By assumption, the solution $g_{\alpha\bar{\beta}}(x, t)$, $(-\infty < t < +\infty)$, has bounded holomorphic bisectional curvature, hence K is finite. Thus, by the derivative estimate of Shi [13] (see also Theorem 7.1 in [12]), the all derivatives of the curvature are also uniformly bounded. In particular, we can use the maximum principle of Cheng–Yau (see Proposition 1.6 in [6]) and then, as observed in [12], this implies that the maximum principle of Hamilton in [9] actually works for the evolution equations of the curvature of $\tilde{g}_{\alpha\bar{\beta}}(x, t)$ on the complete noncompact manifold M . Thus, from [9], we obtain

$$\begin{aligned} \frac{d(a_1)_{\min}}{dt} &\geq ((a_1)_{\min})^2 + 2(a_2)_{\min} \cdot (a_3)_{\min} \\ &\geq 3((a_1)_{\min})^2 \end{aligned}$$

by (2.2). Then, for fixed $t_0 \in (-\infty, 0)$ and $t > t_0$,

$$\begin{aligned} (a_1)_{\min}(t) &\geq \frac{1}{(a_1)_{\min}^{-1}(t_0) - 3(t - t_0)} \\ &\geq \frac{1}{-K^{-1} - 3(t - t_0)}. \end{aligned}$$

Letting $t_0 \rightarrow -\infty$, we get

$$a_1 \geq 0, \quad \text{for all } (x, t) \in M \times (-\infty, 0] \quad (2.3)$$

i.e. A is nonnegative definiteness.

Finally, to prove the nonnegativity of the curvature operator R_m , we recall its corresponding ODE from [9],

$$\frac{dR_m}{dt} = R_m^2 + \begin{pmatrix} 0 & 0 \\ 0 & A^\# \end{pmatrix},$$

where $A^\#$ is the adjoint matrix of A .

Let m_1 be the smallest eigenvalue of the curvature operator R_m . Exactly as above, by using the maximum principle of Hamilton, we have

$$\frac{d(m_1)_{\min}}{dt} \geq (m_1)_{\min}^2$$

where $(m_1)_{\min}(t) = \inf_{x \in M} m_1(x, t)$. Therefore, by the same reasoning in the derivation of (2.3), we have

$$m_1 \geq 0, \quad \text{for all } (x, t) \in M \times (-\infty, +\infty). \quad (2.4)$$

So R_m is nonnegative definite everywhere and the proof of the lemma is completed. *Q.E.C.*

3. Blow Down Argument

This section is devoted to the proof of the main result.

The proof of the theorem.

Let $(M, g_{\alpha\bar{\beta}}(x))$ be a nonflat Kähler-Ricci soliton on a complex two-dimensional Kähler manifold with bounded and nonnegative holomorphic bisectional curvature. Denote $g_{\alpha\bar{\beta}}(x, t), t \in (-\infty, +\infty)$, to be the solution to the Ricci flow (1.1) corresponding to the Kähler-Ricci soliton $g_{\alpha\bar{\beta}}(x)$. The metric $g_{\alpha\bar{\beta}}(x, t)$ is obtained from the pullback of $g_{\alpha\bar{\beta}}(x)$ by a one-parameter group of biholomorphism. Of course, for each $t \in (-\infty, +\infty)$, $g_{\alpha\bar{\beta}}(x, t)$ also has bounded and nonnegative holomorphic bisectional curvature.

We prove by contradiction. Thus, suppose $g_{\alpha\bar{\beta}}(x)$ is of maximal volume growth. This is, there exist some point $x_0 \in M$ and some positive constant C_0 such that

$$\text{Vol}(B(x_0, r)) \geq C_0 r^4 \quad \text{for all } 0 \leq r < +\infty. \quad (3.1)$$

Note that $g_{\alpha\bar{\beta}}(x)$ has nonnegative Ricci curvature. Then it follows from the standard volume comparison that the above inequality actually holds for all $x_0 \in M$. Hence

$$\text{Vol}_t(B_t(x, r)) \geq C_0 r^4 \quad \text{for all } 0 \leq r < +\infty \text{ and } x \in M, \quad (3.2)$$

where $\text{Vol}_t(B_t(x, r))$ denotes the volume of the geodesic ball $B_t(x, r)$ of radius r with center at x with respect to the metric $g_{\alpha\bar{\beta}}(x, t)$.

We first observe that

$$\limsup_{d(x, x_0) \rightarrow +\infty} R(x) d^2(x, x_0) = +\infty \quad (3.3)$$

where $d(x, x_0)$ the geodesic distance between two points $x, x_0 \in M$ with respect to the metric $g_{\alpha\bar{\beta}}(x)$.

In fact, suppose not, thus the curvature of the initial metric $g_{\alpha\bar{\beta}}(x, 0) = g_{\alpha\bar{\beta}}(x)$ in (1.1) has quadratic decay. By applying a result of Shi (see Theorem 8.2 in [14]), we see that

$$\lim_{t \rightarrow +\infty} \sup \{ R(x, t) \mid x \in M \} = 0. \quad (3.4)$$

where $R(x, t)$ is the scalar curvature of the solution $g_{\alpha\bar{\beta}}(x, t)$. But $g_{\alpha\bar{\beta}}(x, t)$ is a nonflat Kähler-Ricci soliton. This is impossible.

With the estimate (3.3), we can then apply a lemma of Hamilton (Lemma 22.2 in [12]) to find a sequence of points x_j , a sequence of radii r_j and a sequence of positive numbers δ_j , $j = 1, 2, \dots$, with $\delta_j \rightarrow 0$ such that

- (a) $R(x, 0) \leq (1 + \delta_j) R(x_j, 0)$ for all x in the ball $B(x_j, r_j)$ of radius r_j centered at x_j with respect to the metric $g_{\alpha\bar{\beta}}$;

- (b) $r_j^2 R(x_j, 0) \rightarrow +\infty$;
- (c) if $s_j = d(x_j, x_0)$, then $\lambda_j = s_j/r_j \rightarrow +\infty$;
- (d) the balls $B(x_j, r_j)$ are disjoint.

We have shown in the previous section that the metric $g_{\alpha\bar{\beta}}$ has nonnegative definite curvature operator. In particular, the sectional curvature is nonnegative. Denote the minimum of the sectional curvature of the metric $g_{\alpha\bar{\beta}}$ at x_j by ν_j . We claim that the following holds

$$\varepsilon_j = \frac{\nu_j}{R(x_j, 0)} \rightarrow 0 \quad \text{as } j \rightarrow +\infty. \quad (3.5)$$

Suppose not, there exists a subsequence $j_k \rightarrow +\infty$ and some positive number $\varepsilon > 0$ such that

$$\varepsilon_{j_k} = \frac{\nu_{j_k}}{R(x_{j_k}, 0)} \geq \varepsilon \quad \text{for all } k = 1, 2, \dots. \quad (3.6)$$

Since the solution $g_{\alpha\bar{\beta}}(x, t)$ exists for $-\infty < t < +\infty$, it follows from the Li-Yau type differential inequality of Cao [1] that the scalar curvature $R(x, t)$ is pointwisely nondecreasing in time. Then, by using the local derivative estimate of Shi [13] (or see Theorem 13.1 in [12]) and (a), (b), we have

$$\begin{aligned} \sup_{x \in B(x_{j_k}, r_{j_k})} |\nabla R_m(x, 0)|^2 &\leq C_1 R^2(x_j, 0) \left(\frac{1}{r_{j_k}^2} + R(x_j, 0) \right) \\ &\leq 2C_1 R^3(x_j, 0), \end{aligned} \quad (3.7)$$

where C_1 is a positive constant depending only on the dimension.

For any $x \in B(x_{j_k}, r_{j_k})$, we obtain from (3.6) and (3.7) that the minimum of the sectional curvature $\nu(x)$ of $g_{\alpha\bar{\beta}}$ at x , satisfies

$$\begin{aligned} \nu(x) &\geq \nu_{j_k} - \sqrt{2C_1} R^{3/2}(x_{j_k}, 0) d(x, x_{j_k}) \\ &\geq R(x_{j_k}, 0) \left(\varepsilon - \sqrt{2C_1} \cdot \sqrt{R(x_{j_k}, 0)} \cdot d(x, x_{j_k}) \right) \\ &\geq \frac{\varepsilon}{2} R(x_{j_k}, 0) \end{aligned} \quad (3.8)$$

if

$$d(x, x_{j_k}) \leq \frac{\varepsilon}{2\sqrt{2C_1} \cdot \sqrt{R(x_{j_k}, 0)}}.$$

Thus, from (a) and (3.8), there exists $k_0 > 0$ such that for any $k \geq k_0$ and

$$x \in B(x_{j_k}, \frac{\varepsilon}{2\sqrt{2C_1} \cdot \sqrt{R(x_{j_k}, 0)}}),$$

we have

$$\frac{\varepsilon}{2}R(x_{j_k}, 0) \leq \text{the sectional curvature at } x \leq 2R(x_{j_k}, 0). \quad (3.9)$$

Therefore the balls $B(x_{j_k}, \frac{\varepsilon}{2\sqrt{2C_1} \cdot \sqrt{R(x_{j_k}, 0)}}), k_0 \leq k < +\infty$, are a family of disjoint remote curvature β -bumps for some $\beta > 0$ in the sense of Hamilton [12]. But this contradicts with the finite bumps theorem of Hamilton [12]. So we have proven the claim (3.5).

Next we blow down the Kähler-Ricci soliton $g_{\alpha\bar{\beta}}(x, t)$ along the points x_j . For the above chosen x_j , r_j and δ_j , let x_j be the new origin, dilate the space by a factor λ_j so that $R(x_j, 0)$ become 1 at the origin at $t = 0$, and dilate in time by λ_j^2 so that it is still a solution to the Ricci flow. The balls $B(x_j, r_j)$ are dilated to the balls centered at the origin of radii $\tilde{r}_j = r_j^2 R(x_j, 0) \rightarrow +\infty$ (by (b)). Since the scalar curvature of $g_{\alpha\bar{\beta}}(x, t)$ is pointwise nondecreasing in time by the Cao's inequality [1], the curvature bounds on $B(x_j, r_j)$ also give bounds for previous times in these balls.

On the other hand, as shown in [4], the combination of the maximal volume growth estimate (3.1) and the local injectivity radius estimate of Cheeger, Gromov and Taylor [3] implies that

$$\text{inj}_M(x_j, g_{\alpha\bar{\beta}}) \geq \frac{\alpha}{\sqrt{R(x_j, 0)}},$$

for some positive constant α independent of j .

So we have everything to take a limit for the dilated solutions. By applying the compactness theorem in [11] and combining (3.2), (3.5), (a) and (b), we

obtain a complete noncompact solution $(\widetilde{M}, \widetilde{g}_{\alpha\bar{\beta}}(x, t))$ of the Ricci flow for $t \in (-\infty, 0]$ such that

- (e) the curvature operator is still nonnegative;
- (f) the scalar curvature $\widetilde{R}(x, t) \leq 1$, for all $x \in \widetilde{M}$, $t \in (-\infty, 0]$, and $\widetilde{R}(0, 0) = 1$;
- (g) the volume of the geodesic ball to metric $\widetilde{g}_{\alpha\bar{\beta}}(x, t)$ still satisfies

$$\text{Vol}_t(\widetilde{B}_t(x, r)) \geq C_0 r^4$$

for all $x \in \widetilde{M}$, $0 \leq r < +\infty$;

- (h) there exists a complex 2-plane at the origin so that at $t = 0$, the corresponding sectional curvature vanishes.

If we consider the universal covering of $\widetilde{M}$, the induced metric of $\widetilde{g}_{\alpha\bar{\beta}}(x, t)$ on the universal covering is clearly still a solution to the Ricci flow and satisfies all of above (e), (f), (g), (h). Thus, without loss of generality, we may assume that $\widetilde{M}$ is simply connected.

By using the strong maximum principle on the evolution equation of the curvature operator of $\widetilde{g}_{\alpha\bar{\beta}}(x, t)$ as in [9] (see Theorem 8.3 of [9]), we know that there exists a constant $K > 0$ such that on the time interval $-\infty < t < -K$, the image of the curvature operator of $(\widetilde{M}, \widetilde{g}_{\alpha\bar{\beta}}(x, t))$ is a fixed Lie subalgebra of $so(4)$ of constant rank on $\widetilde{M}$. Because $\widetilde{M}$ is Kähler, the possibilities are limited to $u(2)$, $so(2) \times so(2)$ or $so(2)$.

In the case $u(2)$, the sectional curvature is strictly positive. Thus, this case is ruled out by (h). In the cases $so(2) \times so(2)$ or $so(2)$, according to [9], the simply connected manifold $\widetilde{M}$ splits as a product $\widetilde{M} = \Sigma_1 \times \Sigma_2$, where Σ_1 and Σ_2 are two Riemann surfaces with nonnegative curvature (by (e)), and at least one of them, say Σ_1 , has positive curvature (by (f)).

Denote by $\tilde{g}_{\alpha\bar{\beta}}^{(1)}(x, t)$ the corresponding metric on Σ_1 . Clearly, it follows from (g) and standard volume comparison that for any $x \in \Sigma_1$, $t \in (-\infty, -K)$, we have

$$\text{Vol}B_{\Sigma_1}(x, r) \geq C_2 r^2 \quad \text{for } 0 \leq r < +\infty \quad (3.10)$$

where both the geodesic ball $B_{\Sigma_1}(x, r)$ and the volume are taken with respect to the metric $\tilde{g}_{\alpha\bar{\beta}}^{(1)}(x, t)$ on Σ_1 , C_2 is a positive constant depending only on C_0 . Also as the curvature of $\tilde{g}_{\alpha\bar{\beta}}^{(1)}(x, t)$ is positive, it follows from Cohn–Vossen inequality that

$$\int_{\Sigma_1} \tilde{R}^{(1)}(x, t) d\sigma_t \leq 8\pi, \quad (3.11)$$

where $\tilde{R}^{(1)}(x, t)$ is the scalar curvature of $(\Sigma_1, \tilde{g}_{\alpha\bar{\beta}}^{(1)}(x, t))$ and $d\sigma_t$ is the volume element of the metric $\tilde{g}_{\alpha\bar{\beta}}^{(1)}(x, t)$.

Now, the metric $\tilde{g}_{\alpha\bar{\beta}}^{(1)}(x, t)$ is a solution to the Ricci flow on the Riemann surface Σ_1 over the time interval $(-\infty, -K)$. The estimates (3.10) and (3.11) imply that for each $t \in (-\infty, -K)$, the curvature of $\tilde{g}_{\alpha\bar{\beta}}^{(1)}(x, t)$ has quadratic decay in the average sense of Shi [14] and then the a priori estimate of Shi (see Theorem 8.2 in [14]) implies that the solution $\tilde{g}_{\alpha\bar{\beta}}^{(1)}(x, t)$ exists for all $t \in (-\infty, +\infty)$ and satisfies

$$\lim_{t \rightarrow +\infty} \sup \left\{ \tilde{R}^{(1)}(x, t) \mid x \in \Sigma_1 \right\} = 0. \quad (3.12)$$

Again, by the inequality of Cao [1], we conclude that

$$\tilde{R}^{(1)}(x, t) \equiv 0 \quad \text{on } \Sigma_1 \times (-\infty, +\infty).$$

This contradicts with the fact that $(\Sigma_1, \tilde{g}_{\alpha\bar{\beta}}^{(1)}(x, t))$ has positive curvature for $t < -K$. Hence we have completed the proof of the theorem. Q.E.C.

References

- [1] Cao, H.D., *On Harnack's inequalities for the Kähler-Ricci flow*, Invent. Math. **109** (1992) 247-263.

- [2] Cao, H.D., *Limits of solutions to the Kähler-Ricci flows*, J. Differential Geometry **45** (1997), 257-272.
- [3] Cheeger, J., Gromov, M. and Taylor, M., *Finite propagation speed, kernel estimates for functions of the Laplace operator, and the geometry of complete Riemannian manifold*, J. Diff. Geom. **17** (1982), 15-53.
- [4] Chen, B.L. and Zhu, X.P., *On complete noncompact Kähler manifolds with positive bisectional curvature*, preprint (2000).
- [5] Chen, B.L., Tang, S.H. and Zhu, X.P., *A uniformization theorem of complete noncompact Kähler surfaces with positive bisectional curvature*, preprint (2001).
- [6] Cheng, S.Y. and Yau, S.T., *On the existence of a complete Kähler metric on non-compact complex manifolds and the regularity of Fefferman's equation*, Comm. Pure Appl. Math., **33** (1980), 507-544.
- [7] Chow, B., *The Ricci flow on the 2-sphere*, J. Differential Geometry, **17** (1991), 325-334.
- [8] Hamilton, R.S., *Three-manifolds with positive Ricci curvature*, J. Differential Geometry, **17**(1982), 255-306.
- [9] Hamilton, R.S., *Four-manifolds with positive curvature operator*, J. Differential Geometry, **24**(1986), 153-179.
- [10] Hamilton, R.S., *The Ricci flow on surfaces*, Contemporary Mathematics, **71** (1988), 237-261.
- [11] Hamilton, R.S., *A compactness property for solutions of the Ricci flow*, Amer. J. Math., **117** (1995), 545-572.

- [12] Hamilton, R.S., *Formation of singularities in the Ricci flow*, surveys in Diff. Geom.Vol.**2**, Boston, International Press 1995,7-136.
- [13] Shi, W.X., *Deforming the metric on complete Riemannian manifolds*, J. Differential Geometry, **30** (1989), 223-301.
- [14] Shi, W.X., *Ricci flow and the uniformization on complete noncompact Kähler manifolds*, J. Differential Geometry, **45** (1997),94-220.